

Alexa Fluor® 488 anti-BTK Phospho (Tyr223) Antibody

Catalog# / Size	601705 / 25 tests 601706 / 100 tests
Clone	A16128B
Regulatory Status	RUO
Other Names	Bruton's tyrosine kinase, Bruton agammaglobulinemia tyrosine kinase, AGMX1, ATK, IMD1, PSCTK1, XLA, BPK
Isotype	Mouse IgG2a, κ
Description	Bruton's tyrosine kinase (BTK) is a Tec family kinase that plays a critical role in B cell development. Upon B cell receptor engagement, BTK translocates to the plasma membrane, where it is transphosphorylated by LYN and SYK kinases at Tyr 551. This initial phosphorylation event is followed by autophosphorylation at Tyr 223. Phosphorylated Tyr 223 is proposed to serve as a docking site for other proteins containing an SH2 domain. Downregulation of BTK activity is achieved through phosphorylation at Ser 180 by PKCβ, which results in reduced membrane recruitment and concomitant transphosphorylation. BTK-mediated B cell receptor signaling is required for B cell survival in the bone marrow and is a therapeutic target for certain B cell malignancies. Mutations in the <i>BTK</i> gene results in X-linked agammaglobulinemia (XLA), a primary immunodeficiency disease characterized by a failure in B cell maturation. Additionally, BTK plays a role in Toll-like receptor signaling in innate immune cells, and has been shown to play a direct role in NLRP3 inflammasome activation, thus contributing to ischaemic brain injury.

Product Details

Verified Reactivity	Human, Mouse
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human BTK peptide phosphorylated at Tyr 223, Complete Freund's adjuvant.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm. Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation. View full statement regarding label licenses
Excitation Laser	Blue Laser (488 nm)

Application Notes

A16128B was generated in the same immunization that produced clone A16128C (anti-BTK Phospho (Tyr223)/ITK Phospho (Tyr180)). Both clones bind BTK Phospho Tyr223 with similar affinity, but in comparison to clone A16128C, clone A16128B has weaker recognition of phosphorylated Tyrosine 180 of ITK by intracellular staining and flow cytometric analysis.

Antigen Details

Structure	BTK encodes a 659 amino acid protein with a predicted molecular weight of 76.2 kD. Contains a PH domain (a.a. 1-133) that binds phosphatidylinositol (3,4,5)-triphosphate (PIP3) and polyphosphates, an SH3 domain (a.a. 214-274), an SH2 domain (a.a. 281-377), and a protein kinase domain (402-655).
Distribution	Cytosol, plasma-membrane associated
Function	Links B cell receptor engagement with downstream signaling events
Interaction	ARID3A, BLNK, CAV1, GNAQ, GTF2I, PLGC2, PRKD1, SH3BP5
Ligand/Receptor	PIP3
Biology Area	Cell Biology, Signal Transduction
Molecular Family	Phospho-Proteins, Protein Kinases/Phosphatase
Antigen References	1. Krysiak K, <i>et al.</i> 2017. <i>Blood</i>. 129:473. 2. Burger JA, <i>et al.</i> 2016. <i>Nat Commun</i>. 7:11589. 3. Wang Q, <i>et al.</i> 2015. <i>Elife</i>. 4:e06074. 4. Ito M, <i>et al.</i> 2015. <i>Nat. Commun</i>. 6:7360. 5. Pillinger G, <i>et al.</i> 2015. <i>Sci. Rep</i>. 5:12949. 6. Cinar M, <i>et al.</i> 2013. <i>Leuk. Res</i>. 37:1271. 7. Doyle SL, <i>et al.</i> 2007 <i>J. Biol. Chem</i>. 282:36953. 8. Nore BF, <i>et al.</i> 2003. <i>Biochem. Biophys. Acta</i>. 1645:123. 9. Li Z, <i>et al.</i> 1997. <i>Proc. Natl. Acad. Sci</i>. 94:13820.

Gene ID [695](#)


Related Protocols

- [Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-BTK Phospho (Tyr223), Alexa Fluor® 488 anti-BTK Phospho (Tyr223), Purified anti-BTK Phospho (Tyr 223), Alexa Fluor® 647 anti-BTK Phospho (Tyr223) Antibody

Product Data



PBMC untreated (middle panel) or treated with H₂O₂ (right panel) to induce phosphorylation of BTK were fixed and permeabilized using the True-Phos™ Perm Buffer Set (Cat. No. 425401) and intracellularly stained with Alexa Fluor® 488 anti-BTK Phospho (Tyr223) (clone A16128B) (middle and right panels) or Alexa Fluor® 488 Mouse IgG2a, κ isotype Control (left panel, representative plot for untreated and H₂O₂-treated cells) (Cat. No. 400233). Cells were co-stained with APC anti-CD20 (cytoplasmic) (Cat. No. 340516).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587