

APC anti-human CD11b Antibody

Catalog# / Size	340007 / 25 tests 340008 / 100 tests
Clone	W18347A
Regulatory Status	RUO
Other Names	Integrin α M chain, C3biR, CR3, Mac-1, Mo1, ITGAM
Isotype	Rat IgG2a, κ
Description	CD11b is a 165-170 kD type I transmembrane glycoprotein also known as α M integrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β 2 (CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils. It is also involved in granulocyte adhesion, phagocytosis, and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen, and factor X.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Human CD11b recombinant protein
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)
RRID	AB_3662236 (BioLegend Cat. No. 340007) AB_3662236 (BioLegend Cat. No. 340008)

Antigen Details

Structure	Integrin, type I transmembrane glycoprotein, associates with integrin β 2 (CD18), 165-170 kD
Distribution	Granulocytes, monocytes/macrophages, dendritic cells, NK cells, subset of T cells, subset of B cells
Function	Adhesion, phagocytosis, chemotaxis, neutrophil activation

Interaction	B cells, Dendritic cells, Granulocytes, Macrophages, Monocytes, Neutrophils, NK cells, T cells, Tregs
Ligand/Receptor	ICAM-1(CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen, factor X
Cell Type	B cells, Dendritic cells, Granulocytes, Monocytes, Neutrophils, NK cells, T cells, Tregs
Biology Area	Cell Adhesion, Cell Biology, Immunology, Innate Immunity
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	1. Stewart M, <i>et al.</i> 1995. <i>Curr. Opin. Cell Biol.</i> 7:690.
Gene ID	3684


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD11b Antibody, APC/Fire™ 750 anti-human CD11b, PE anti-human CD11b, PE/Cyanine7 anti-human CD11b, APC anti-human CD11b, PerCP/Cyanine5.5 anti-human CD11b, Brilliant Violet 750™ anti-human CD11b Antibody

Product Data



Human peripheral blood mononuclear cells were stained with anti-human CD14 (clone HCD14) Brilliant Violet 421™ and anti-human CD11b (clone W18347A) APC (left) or rat IgG2a, κ APC isotype control (right). Data shown was gated on monocytes.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587