

APC anti-human IL-1RL1 (ST2) Antibody

Catalog# / Size	379411 / 25 tests 379412 / 100 tests
Clone	A16008J
Regulatory Status	RUO
Other Names	IL-33R, ST2, IL-1RL1
Isotype	Mouse IgG2b, κ
Description	IL1RL1 is a member of the Toll-like receptor superfamily the receptor for Interleukin-33. It is a 556 amino acid single-pass type I membrane protein. ST2 exists in two forms as splice variants: a soluble form, which acts as a decoy receptor and sequesters free IL-33, and a membrane-bound form (ST2), which activates the MyD88/NF- κ B signaling pathway. Signaling through IL1RL1 drives cytokine production in mast cells, basophils, eosinophils, invariant NKT and NK cells, Th2 lymphocytes, and type 2 innate immune cells. IL-33 signaling is required for maintaining the suppressive function of regulatory T cells.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Recombinant human IL-1RL1, AA19-328
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)

Antigen Details

Structure	Single-pass type I membrane protein
Distribution	Th2 lymphocytes, mast cells, eosinophils, basophils, smooth muscle cells, endothelial cells
Function	Cytokine signaling
Interaction	Signaling requires association of the coreceptor IL1RAP. Recruits MYD88, IRAK1, IRAK4, and TRAF6.

Ligand/Receptor	IL-33
Cell Type	Basophils, Endothelial cells, Eosinophils, Lymphocytes, NK cells, Th2, Tregs
Biology Area	Immunology
Molecular Family	Cytokine/Chemokine Receptors
Antigen References	1. Schmitz J, <i>et al.</i> 2005. <i>Immunity</i>. 23: 479-490. 2. Brint E, <i>et al.</i> 2004. <i>Nat Immun.</i> 5:373-379. 3. Griesenauer B, <i>et al.</i> 2017. <i>Front Immunol.</i> 8:475.
Gene ID	9173


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

PE/Dazzle™ 594 anti-human IL-1RL1 (ST2), Purified anti-human IL-1RL1 (ST2), PE anti-human IL-1RL1 (ST2), PE/Cyanine7 anti-human IL-1RL1 (ST2), APC anti-human IL-1RL1 (ST2), Brilliant Violet 421™ anti-human IL-1RL1 (ST2)

Product Data



IL-3 stimulated (overnight) human peripheral blood mononuclear cells were stained with anti-human HLA-DR (clone L243) Brilliant Violet 421™, anti-human CD123 (clone S18016C) PerCP/Fire™ 806, and anti-human IL-1RL1 (ST2) (clone A16008J) APC (left) or mouse IgG2b, κ APC isotype control (right). Data shown was gated on HLA-DR negative population.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587