

APC anti-mouse CD69 Recombinant Antibody

Catalog# / Size	164205 / 25 µg 164206 / 100 µg
Clone	QA17A41
Regulatory Status	RUO
Other Names	Very Early Activation Antigen (VEA), AIM, EA1, MLR3, gp34/28
Isotype	Rat IgG1, κ
Description	CD69 is a 60 kD type II membrane protein composed of a 27/33 kD disulfide-linked homodimer, also known as Very Early Activation (VEA) antigen, AIM, EA1, MLR3, and gp34/28. It is expressed on a subset of thymocytes and platelets. CD69 is rapidly induced on activated T and B cells, neutrophils, and NK cells. It is a C-type lectin, closely related to the NKR-P1 and Ly-49 NK cell activation molecules. CD69 is involved in the early events of cell activation and thymocyte positive selection.

Product Details

Verified Reactivity	Mouse
Antibody Type	Recombinant
Host Species	Rat
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	0.2 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)
RRID	AB_3698929 (BioLegend Cat. No. 164205)

Antigen Details

Structure	C-type lectin, 27/33 kD
Distribution	Activated T cells and B cells, NK cells, granulocytes, thymocytes, platelets
Function	Lymphocyte activation
Cell Type	B cells, Granulocytes, NK cells, Platelets, T cells, Thymocytes, Tregs
Biology Area	Costimulatory Molecules, Immunology, Innate Immunity
Molecular Family	CD Molecules


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-mouse CD69 Recombinant Antibody, PE anti-mouse CD69 Recombinant Antibody, APC anti-mouse CD69 Recombinant Antibody

Product Data



PMA+ionomycin stimulated (6 hours) C57BL/6 mouse splenocytes were stained with anti-mouse CD69 recombinant (clone QA17A41) APC (filled histogram) or rat IgG1, κ APC isotype control (open histogram).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587