

APC/Fire™ 750 anti-human CD4 Antibody

Catalog# / Size	317461 / 25 tests 317462 / 100 tests
Clone	OKT4
Regulatory Status	RUO
Workshop	HCDM listed
Other Names	T4
Isotype	Mouse IgG2b, κ
Description	CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with IL-16.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Reported Reactivity	Chimpanzee
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human peripheral T cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 774 nm.
Excitation Laser	Red Laser (633 nm)
Application Notes	The OKT4 antibody binds to the D3 domain of CD4 and does not block HIV binding. Additional reported applications (for the relevant formats) include: immunohistochemistry of frozen sections and blocking of T cell activation. This clone was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 317453 and 317454). In a small subset of individuals, the OKT4 clone does not bind to CD4 due to polymorphisms in CD4. ⁹

Application References

(PubMed link indicates
BioLegend citation)

1. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
2. Reinherz EL, *et al.* 1979. *Proc. Natl. Acad. Sci.* 76:4061.
3. Kmiecik M, *et al.* 2009. *J. Transl. Med.* 7:89. (FC) [PubMed](#)
4. Cicin-Sain L, *et al.* 2010. *J. Immunol.* 184:6739. [PubMed](#)
5. Rosenzweig M, *et al.* 2001. *J. Med. Primatol.* 30:36.
6. Linder J, *et al.* 1987. *Am. J. Pathol.* 127:1.
7. Boche D, *et al.* 1999. *J. Neurovirol.* 5:232. (IHC)
8. Reinherz EL, *et al.* 1979. *Proc. Natl. Acad. Sci. USA.* 76:4061. (Immunogen)
9. Lederman S, *et al.* 1991. *Mol Immunol.* 28:1171-81.

RRID

AB_3683334 (BioLegend Cat. No. 317461)
AB_3683334 (BioLegend Cat. No. 317462)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, 55 kD
Distribution	T cell subset, majority of thymocytes, monocytes/macrophages
Function	MHC class II co-receptor, lymphocyte adhesion, thymic differentiation, HIV receptor
Ligand/Receptor	MHC class II molecules, HIV gp120, IL-16
Cell Type	Macrophages, Monocytes, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	1. Center D, <i>et al.</i> 1996. <i>Immunol. Today</i> 17:476. 2. Gaubin M, <i>et al.</i> 1996. <i>Eur. J. Clin. Chem. Clin. Biochem.</i> 34:723.
Gene ID	920


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Brilliant Violet 650™ anti-human CD4, Purified anti-human CD4, Biotin anti-human CD4, FITC anti-human CD4, PE anti-human CD4, PE/Cyanine5 anti-human CD4, PE/Cyanine7 anti-human CD4, APC anti-human CD4, APC/Cyanine7 anti-human CD4, Alexa Fluor® 488 anti-human CD4, Alexa Fluor® 647 anti-human CD4, Alexa Fluor® 700 anti-human CD4, Pacific Blue™ anti-human CD4, PerCP/Cyanine5.5 anti-human CD4, PerCP anti-human CD4, Brilliant Violet 421™ anti-human CD4, Brilliant Violet 605™ anti-human CD4, Brilliant Violet 711™ anti-human CD4, Brilliant Violet 785™ anti-human CD4, Brilliant Violet 510™ anti-human CD4, Brilliant Violet 570™ anti-human CD4, PE/Dazzle™ 594 anti-human CD4, TotalSeq™-A0922 anti-human CD4, Ultra-LEAF™ Purified anti-human CD4, TotalSeq™-C0922 anti-human CD4, TotalSeq™-B0922 anti-human CD4, PE/Fire™ 810 anti-human CD4, APC/Fire™ 750 anti-human CD4 Antibody

Product Data



Human peripheral blood lymphocytes were stained with anti-human CD3 (clone UCHT1) FITC and anti-human CD4 (clone OKT4) APC/Fire™ 750 (left) or mouse IgG2b, κ APC/Fire™ 750 isotype control (right)

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587