

Brilliant Violet 570™ anti-human CD326 (Ep-CAM) Antibody

Catalog# / Size	324261 / 25 tests 324262 / 100 tests
Clone	9C4
Regulatory Status	RUO
Other Names	Ep-CAM, tumor associated calcium signal transducer 1, epithelial cell surface antigen, epithelial glycoprotein 2, EGP2, adenocarcinoma associated antigen, TROP1
Isotype	Mouse IgG2b, κ
Description	CD326 is also known as Ep-CAM, tumor associated calcium signal transducer 1, epithelial cell surface antigen, epithelial glycoprotein 2, EGP2, adenocarcinoma associated antigen, and TROP1. CD326 is a type I transmembrane protein containing six disulfide bridges and one THYRO domain. This cell surface glycosylated 40 kD protein is highly expressed in bone marrow, colon, lung, and most normal epithelial cells and is expressed on carcinomas of gastrointestinal origin. Recently, it has been reported that CD326 expression occurs during the early steps of erythropoiesis. CD326 functions as a homotypic calcium-independent cell adhesion molecule and is believed to be involved in carcinogenesis by its ability to induce genes involved in cellular metabolism and proliferation. CD326 antigen is an immunotherapeutic target for the treatment of human carcinomas.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	DU.4475 breast carcinoma
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 570™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Brilliant Violet 570™ excites at 405 nm and emits at 570 nm. The bandpass filter 585/42 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 570™ is a trademark of Sirigen Group Ltd.

[Learn more about Brilliant Violet™.](#)

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in

any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Excitation Laser Violet Laser (405 nm)

Application Notes Additional reported applications (for the relevant formats) include: immunofluorescence, immunohistochemistry³, and spatial biology (IBEX)^{4,5}.

Application References

(PubMed link indicates BioLegend citation)

1. Lammers R, *et al.* 2002. *Exp. Hematol.* 30:537.
2. Schultz LD, *et al.* 2010. *P. Natl. Acad. Sci. USA* 107:13022. [PubMed](#)
3. Human Protein Atlas <http://www.proteinatlas.org/ENSG00000119888/antibody> (IHC)
4. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci USA*. 117:33455-33465. (SB) [PubMed](#)
5. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

RRID AB_3720529 (BioLegend Cat. No. 324261)
AB_3720529 (BioLegend Cat. No. 324262)

Antigen Details

Structure	Type I transmembrane protein, contains six disulfide bridges, one THYRO domain, approximate molecular weight 40 kD.
Distribution	Highly expressed in bone marrow, colon, lung, and most normal epithelial cells. Also highly expressed on carcinomas of gastrointestinal origin. Expressed during early erythropoiesis.
Function	Homotypic calcium-independent cell adhesion. CD326 is believed to be involved in carcinogenesis by its ability to induce genes involved in cellular metabolism and proliferation.
Modification	Glycosylated.
Cell Type	Embryonic Stem Cells, Epithelial cells
Biology Area	Cell Biology, Immunology, Stem Cells
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	1. Strnad J, <i>et al.</i> 1989. <i>Cancer Res.</i> 49:314. 2. Munz M, <i>et al.</i> 2004. <i>Oncogene</i> 23:5748. 3. Rao CG, <i>et al.</i> 2005. <i>Int. J. Oncol.</i> 27:49.
Gene ID	4072


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

PE/Cyanine5 anti-human CD326 (EpCAM), TotalSeq™-D0123 anti-human CD326 (EpCAM), Spark UV™ 387 anti-human CD326 (EpCAM), Spark Red™ 718 anti-human CD326 (EpCAM) (Flexi-Fluor™), PE/Dazzle™ 594 anti-human CD326 (EpCAM), Brilliant Violet 510™ anti-human CD326 (EpCAM), Brilliant Violet 785™ anti-human CD326 (EpCAM), APC/Fire™ 750 anti-human CD326 (EpCAM), Biotin anti-human CD326 (EpCAM), Pacific Blue™ anti-human CD326 (EpCAM), Brilliant Violet 605™ anti-human CD326 (EpCAM), Brilliant Violet 421™ anti-human CD326 (EpCAM), Brilliant Violet 650™ anti-human CD326 (EpCAM), Alexa Fluor® 594 anti-human CD326 (EpCAM), Brilliant Violet 750™ anti-human CD326 (EpCAM), Purified anti-human CD326 (EpCAM) (Maxpar® Ready), Spark YG™ 581 anti-human CD326 (EpCAM) (Flexi-Fluor™), Spark YG™ 593 anti-human CD326 (EpCAM) (Flexi-Fluor™), Alexa Fluor® 647 anti-human CD326 (EpCAM), PerCP/Cyanine5.5 anti-human CD326 (EpCAM), Brilliant Violet 711™ anti-human CD326 (EpCAM), TotalSeq™-C0123 anti-human CD326 (EpCAM), TotalSeq™-B0123 anti-human CD326 (EpCAM), Spark PLUS UV395™ anti-human CD326 (EpCAM), PE anti-human CD326, Brilliant Violet 570™ anti-human CD326 (EpCAM), Spark NIR™ 685 anti-human CD326 (EpCAM) (Flexi-Fluor™), TotalSeq™-A0123 anti-human CD326 (EpCAM), Alexa Fluor® 700 anti-human CD326 (EpCAM), PE/Cyanine7 anti-human CD326 (EpCAM), Spark Blue™ 574 anti-human CD326 (EpCAM) (Flexi-Fluor™), Spark Blue™ 550 anti-human CD326 (EpCAM) (Flexi-Fluor™), Purified anti-human CD326 (EpCAM), FITC anti-human CD326 (EpCAM), PE anti-human CD326 (EpCAM), APC anti-human CD326 (EpCAM), Alexa Fluor® 488 anti-human CD326 (EpCAM), APC/Cyanine7 anti-human CD326 (EpCAM)

Product Data



HT-29 cells (human colorectal adenocarcinoma cell line) were stained with anti-human CD326 (Ep-CAM) (clone 9C4) Brilliant Violet 570™ (filled histogram) or mouse IgG2b, κ Brilliant Violet 570™ isotype control (open histogram).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587