

Brilliant Violet 605™ anti-human CD56 (NCAM) Antibody

Catalog# / Size	343761 / 25 tests 343762 / 100 tests
Clone	W22097A
Regulatory Status	RUO
Other Names	Leu-19, NKH1, NCAM-1
Isotype	Rat IgG2a, κ
Description	CD56, also known as NCAM (neural cell adhesion molecule), Leu-19, or NKH1, is a single transmembrane glycoprotein. It is a member of the Ig superfamily. The 140 kD isoform is expressed on NK and NKT cells. CD56 is also expressed in the brain (cerebellum and cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparan sulfate.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Mammalian-derived recombinant protein aa 20-718
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.</p> <p>Learn more about Brilliant Violet™.</p> <p>This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p>
Excitation Laser	Violet Laser (405 nm)

Application Notes

Flow cytometry testing showed that clone W22097A can block the binding of clone HCD56 and W22098E on target cells.

Flow cytometry testing showed that clone W22097A does not affect the binding of clone MEM-188, 5.1H11 and W22098A on target cells.

RRID

AB_3683362 (BioLegend Cat. No. 343761)

AB_3683362 (BioLegend Cat. No. 343762)

Antigen Details

Structure Ig superfamily, single transmembrane or GPI-anchored glycoprotein

Distribution NK cells, T subset, neural tissue, some LGL and myeloid leukemias

Function Adhesion

Ligand/Receptor Heparan sulfate

Cell Type Leukemia, Neurons, NK cells, NKT cells

Biology Area Cell Adhesion, Cell Biology, Costimulatory Molecules, Immunology

Molecular Family Adhesion Molecules, CD Molecules

Antigen References

1. Lanier L, *et al.* 1991. *J Immunol.* 146:4421-6.
2. Hemperly J, *et al.* 1990. *J Mol Neurosci.* 2:71-8.
3. Cremer H, *et al.* 1994. *Nature.* 367:455-9.

Gene ID

[4684](#)


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD56 (NCAM), PE anti-human CD56 (NCAM), PE/Cyanine7 anti-human CD56 (NCAM), APC anti-human CD56 (NCAM) Antibody, Brilliant Violet 421™ anti-human CD56 (NCAM), Brilliant Violet 605™ anti-human CD56 (NCAM)

Product Data



Human peripheral blood lymphocytes were stained with anti-human CD16 (clone 3G8) FITC and anti-human CD56 (NCAM) (clone W22097A) Brilliant Violet 605™ (left) or rat IgG2a, κ Brilliant Violet 605™ isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form,

used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587