

PE anti-human CD28 Recombinant Antibody

Catalog# / Size	377305 / 25 tests 377306 / 100 tests
Clone	QA17A32
Regulatory Status	RUO
Workshop	V-CD28.05
Other Names	T44, Tp44
Isotype	Mouse IgG1, κ
Description	CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. In vitro studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell activation and proliferation.

Product Details

Verified Reactivity	Human
Antibody Type	Recombinant
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Clone QA17A32 completely blocks binding of parental clone. Clone QA17A32 activates human T cells in co-stimulation assay with CD3 as efficiently as the parental clone.

Application References

(PubMed link indicates BioLegend citation)

1. Schlossman S, et al. Eds. 1995. *Leucocyte Typing V* Oxford University Press New York.

2. Nunes J, et al. 1993. *Biochem J.* 293:835.

3. Calea-Lauri J, et al. 1999. *J Immunol.* 163:62.

4. Tazi A, et al. 1999. *J Immunol.* 163:3511. (IHC)

5. Marti F, et al. 2001. *J Immunol.* 166:197. (Costim)

6. Jeong SH, et al. 2004. *J Virol.* 78:6995. (Costim)

7. Rivollier A, et al. 2004. *Blood.* 104:4029. (Costim)

8. Scharschmidt E, et al. 2004. *Mol Cell Biol.* 24:3860. (Costim)

9. Sheng W, et al. 2007. *Elsevier.* 580:6819. [PubMed](#)

10. Mitsuhashi M. 2007. *Clin Chem.* 53:148. [PubMed](#)

11. Ye Z, et al. 2008. *Infect Immun.* 76:2541. [PubMed](#)

12. Magatti M, et al. 2008. *Stem Cells.* 26:182. (FA) [PubMed](#)

[See More](#)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, homodimer, 44 kD
Distribution	Mature T cells, thymocytes, NK cell subsets, plasma cells, EBV-positive B cells
Function	T cell costimulation
Interaction	B cells, NK cells, Plasma cells, T cells, Thymocytes, Tregs
Ligand/Receptor	CD80, CD86
Cell Type	T cells, Thymocytes
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules

Antigen References

1. Schlossman S, et al. Eds. 1995. *Leucocyte Typing V* Oxford University Press New York.

2. June CH, et al. 1994. *Immunol Today.* 15:321-31.

3. Linskey PS, et al. 1993. *Annu Rev Immunol.* 11:191-212.

Gene ID [940](#)


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Ultra-LEAF™ Purified anti-human CD28 Recombinant Antibody, PE anti-human CD28 Recombinant Antibody

Product Data



Human peripheral blood lymphocytes were stained with antihuman CD3 (clone SK7) APC and anti-human CD28 recombinant (clone QA17A32) PE (left) or mouse IgG1, κ PE isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587