

PE anti-mouse CD7 Antibody

Catalog# / Size	167103 / 25 µg 167104 / 100 µg
Clone	S23010B
Regulatory Status	RUO
Isotype	Rat IgG1, κ
Description	CD7, also known as Leu-9, is an approximately 40 kD glycosylated and palmitoylated transmembrane protein in the immunoglobulin superfamily. The extracellular domain of mature mouse CD7 shares 43% and 81% amino acid sequence similarity with human and rat CD7, respectively. Mouse CD7 mRNA has been found in thymus, spleen, bone marrow, and small intestine. Both the ligand and function of mouse CD7 are not yet clear.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Mouse CD7-transfected cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Preparation	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
Concentration	0.2 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
RRID	AB_3698935 (BioLegend Cat. No. 167103) AB_3698935 (BioLegend Cat. No. 167104)

Antigen Details

Structure	Type I transmembrane glycoprotein; Ig superfamily
Distribution	Mouse CD7 mRNA exists in thymus, spleen, bone marrow, and small intestine
Cell Type	Hematopoietic stem and progenitors, Leukemia, NK cells, T cells, Thymocytes
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules

Antigen References

1. Yoshikawa K, *et al.* 1993. *Immunogenetics*. 37:114-9.
2. Haynes BF, *et al.* 1999. *Crit Rev Immunol*. 19:331.
3. Bonilla FA, *et al.* 1997. *Int Immunol*. 9:1875.

Gene ID [12516](#)


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-mouse CD7, PE anti-mouse CD7, Brilliant Violet 421™ anti-mouse CD7, Biotin anti-mouse CD7

Product Data



C57BL/6 mouse splenocytes were stained with anti-mouse CD3 (clone 17A2) FITC and anti-mouse CD7 (clone S23010B) PE (left) or rat IgG1, κ PE isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587