

PE/Cyanine7 anti-human IL-2 Antibody

Catalog# / Size	310613 / 25 tests 310614 / 100 tests
Clone	W19046A
Regulatory Status	RUO
Other Names	Interleukin-2, T cell growth factor (TCGF); Eosinophil differentiation factor (EDF); Killer cell helper factor (KHF); Macrophage-activating factor for cytotoxicity I (MAF-C I); Thymocyte differentiation factor (TDF)
Isotype	Rat IgG2b, κ
Description	IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Recombinant human IL-2
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular flow cytometry using our Cyto-Fast™ Fix/Perm Buffer Set. For flow cytometric staining, the suggested use of this reagent is 5 μ L per 10^6 cells in 100 μ L volume. It is highly recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
RRID	AB_3683322 (BioLegend Cat. No. 310613) AB_3683322 (BioLegend Cat. No. 310614)

Antigen Details

Structure	Cytokine; 15.4 kD (Mammalian)
Distribution	T cells
Function	Upregulated by NFAT; downregulated by TCF-8 and CIF (colostrums inhibitory factor)
Ligand/Receptor	High affinity heterotrimer of IL-2R α / β / γ , intermediate affinity homodimer IL-2R α (CD25; p55; Tac) and heterodimer IL-2R β (CD122)/ γ ; γ -subunit (CD132) in common with IL-4R, IL-7R, IL-13R, IL-

Molecular Family Cytokines/Chemokines

Antigen References

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.
2. Taniguchi T, *et al.* 1993. *Cell* 73:5.
3. Nistico G. 1993. *Prog. Neurobiol.* 40:463.
4. Waldmann T, *et al.* 1993. *Ann. NY Acad. Sci.* 685:603.

Gene ID [3558](#)


Related Protocols

- [Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human IL-2, Spark Violet™ 423 anti-human IL-2, APC anti-human IL-2, FITC anti-human IL-2, PE anti-human IL-2 Antibody, PE/Dazzle™ 594 anti-human IL-2 Antibody, PE/Cyanine7 anti-human IL-2 Antibody

Product Data



Human peripheral blood lymphocytes were stimulated with Cell Activation Cocktail (with Brefeldin A) (Cat. No. 423303) for 6 hours and surface stained with anti-human CD3 (clone UCHT1) Brilliant Violet 421™. Cells were then treated with Cyto-Fast™ Fix/Perm Buffer Set (Cat. No. 426803) and intracellularly stained with anti-human IL-2 (clone W19046A) PE/Cyanine7 (left) or rat IgG2b, κ PE/Cyanine7 isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587