

PE/Fire™ 744 anti-human CD14 Antibody

Catalog# / Size	367165 / 25 tests 367166 / 100 tests
Clone	63D3
Regulatory Status	RUO
Other Names	Monocyte differentiation antigen CD14, myeloid cell-specific leucine-rich glycoprotein, LPS receptor
Isotype	Mouse IgG1, κ
Description	CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein that is also known as the LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens and in the upregulation of adhesion molecules and cytokine expression in monocytes and neutrophils.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Purified human peripheral blood monocytes.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Fire™ 744 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. True-Stain Monocyte Blocker™ (Cat No. 426102) is recommended to minimize non-specific staining of PE/Fire™ 744 on monocytes and macrophages. * PE/Fire™ 744 has a maximum excitation of 565 nm and a maximum emission of 744 nm.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application References	1. Fridlender ZG, <i>et al.</i> 1999. <i>Hum. Immunol.</i> 11:1028. 2. Devitt A, <i>et al.</i> 1998. <i>Nature</i> 6675:505.
(PubMed link indicates BioLegend citation)	
RRID	AB_3720556 (BioLegend Cat. No. 367165)

Antigen Details

Structure	GPI-linked membrane glycoprotein.
Distribution	Monocytes, macrophages, dendritic cells, and granulocytes.
Function	LPS receptor, clearance of Gram-negative pathogens.
Interaction	LPS.
Ligand/Receptor	LPS.
Cell Type	Dendritic cells, Granulocytes, Macrophages, Monocytes, Neutrophils
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	1. Stocks SC, <i>et al.</i> 1990. <i>Biochem. J.</i> 268:275. 2. Wright SD, <i>et al.</i> 1990. <i>Science</i> 4975:1431.
Gene ID	929


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Spark Red™ 718 anti-human CD14 (Flexi-Fluor™), Purified anti-human CD14, PE anti-human CD14, Biotin anti-human CD14, PerCP/Cyanine5.5 anti-human CD14, APC/Fire™ 750 anti-human CD14, APC/Cyanine7 anti-human CD14, Alexa Fluor® 700 anti-human CD14, FITC anti-human CD14, Brilliant Violet 605™ anti-human CD14, Alexa Fluor® 647 anti-human CD14, Alexa Fluor® 488 anti-human CD14, Brilliant Violet 510™ anti-human CD14, Spark YG™ 581 anti-human CD14, Spark Blue™ 574 anti-human CD14, PE/Fire™ 744 anti-human CD14, PE/Cyanine7 anti-human CD14, APC anti-human CD14, Brilliant Violet 750™ anti-human CD14, Spark Blue™ 550 anti-human CD14, Spark NIR™ 685 anti-human CD14, Pacific Blue™ anti-human CD14, APC/Fire™ 810 anti-human CD14, Brilliant Violet 650™ anti-human CD14, PE/Dazzle™ 594 anti-human CD14, TotalSeq™-C0051 anti-human CD14, Brilliant Violet 421™ anti-human CD14, Brilliant Violet 711™ anti-human CD14, TotalSeq™-B0051 anti-human CD14, PE/Fire™ 640 anti-human CD14, PerCP anti-human CD14, TotalSeq™-A0051 anti-human CD14, Brilliant Violet 785™ anti-human CD14, PE/Fire™ 700 anti-human CD14

Product Data



Human peripheral blood monocytes were stained with anti-human CD14 (clone 63D3) PE/Fire™ 744 (filled histogram) or mouse IgG1, κ PE/Fire™ 744 isotype control (open histogram).

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587