

PE/Fire™ 744 anti-mouse I-A/I-E Antibody

Catalog# / Size	107689 / 25 µg 107690 / 100 µg
Clone	M5/114.15.2
Regulatory Status	RUO
Other Names	MHC class II
Isotype	Rat IgG2b, κ
Description	These class II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2 ^{b,d,q,r} bearing mice and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Activated C57BL/6 mouse spleen cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Fire™ 744 under optimal conditions.
Concentration	0.2 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.03 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application. True-Stain Monocyte Blocker™ (Cat No. 426102) is recommended to minimize non-specific staining of PE/Fire™ 744 on monocytes and macrophages. * PE/Fire™ 744 has a maximum excitation of 565 nm and a maximum emission of 744 nm.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	The M5/114.15.2 antibody reacts with a polymorphic determinant shared by the I-A^b, I-A^d, I-A^q, I-E^d, and I-E^k MHC class II alloantigens from mice carrying H-2^{p,r,q,b,d,u} haplotypes. Clone M5/114.15.2 however does not react with I-A^f, I-A^k, or I-A^s MHC class II alloantigens.¹ Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunohistochemistry of frozen sections^{2,3,6}, <i>in vitro</i> and <i>in vivo</i> blocking of antigen presentation or ligand binding⁴⁻⁷, and spatial biology (IBEX)^{17,18}. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. Nos. 107655 & 107656).

Application References

(PubMed link indicates
BioLegend citation)

1. Bhattacharya A, *et al.* 1981. *J. Immunol.* 127:2488. (IP)
2. Viville S, *et al.* 1993. *Cell* 72:635. (IHC)
3. Nelson AJ, *et al.* 1993. *J. Immunol.* 151:2453. (IHC)
4. Shi Y, *et al.* 1998. *J. Exp. Med.* 187:367. (Block)
5. Yamashita I, *et al.* 1993. *Int. Immunol.* 5:1139.
6. Guo M, *et al.* 1995. *Zygote* 3:65. (IHC)
7. Kim A, *et al.* 2004. *Exp. Mol. Med.* 36:428. (Block)
8. Luckashenak NA, *et al.* 2006. *J. Immunol.* 177:5177.
9. Venanzi ES, *et al.* 2007. *J. Immunol.* 179:5693.
10. Christensen SR, *et al.* 2006. *Immunity* 25:417. [PubMed](#)
11. Matte-Martone C, *et al.* 2008. *Blood* 111:3884. [PubMed](#)
12. De Pascalis R, *et al.* 2008. *Infect. Immun.* 76:4311. [PubMed](#)

[See More](#)

Antigen Details

Structure	MHC class II
Distribution	B cell and activated T cells, APCs of the H-2 ^{b,d,q,r} bearing mice
Function	Antigen presentation
Ligand/Receptor	CD3/TCR, CD4
Cell Type	Antigen-presenting cells, B cells, Dendritic cells, T cells, Tregs
Biology Area	Immunology, Innate Immunity
Molecular Family	MHC Antigens
Antigen References	1. Watts C. 1997. <i>Ann. Rev. Immunol.</i> 15:821. 2. Pamer E, <i>et al.</i> 1998. <i>Ann. Rev. Immunol.</i> 16:323.
Gene ID	14961 14969


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Spark YG™ 581 anti-mouse I-A/I-E, PE/Fire™ 810 anti-mouse I-A/I-E, PerCP/Fire™ 806 anti-mouse I-A/I-E, APC/Fire™ 810 anti-mouse I-A/I-E, Purified anti-mouse I-A/I-E (Maxpar® Ready), Brilliant Violet 605™ anti-mouse I-A/I-E, Brilliant Violet 711™ anti-mouse I-A/I-E, PE/Dazzle™ 594 anti-mouse I-A/I-E, APC/Fire™ 750 anti-mouse I-A/I-E, Brilliant Violet 650™ anti-mouse I-A/I-E, Brilliant Violet 785™ anti-mouse I-A/I-E, Alexa Fluor® 594 anti-mouse I-A/I-E, APC/Cyanine7 anti-mouse I-A/I-E, Brilliant Violet 421™ anti-mouse I-A/I-E, Brilliant Violet 510™ anti-mouse I-A/I-E, Spark UV™ 387 anti-mouse I-A/I-E, Spark PLUS UV395™ anti-mouse I-A/I-E, Spark Violet™ 538 anti-mouse I-A/I-E, Brilliant Violet 750™ anti-mouse I-A/I-E, Spark Red™ 718 anti-mouse I-A/I-E, Spark NIR™ 685 anti-mouse I-A/I-E (Flexi-Fluor™) Antibody, StarBright UltraViolet 795 anti-mouse I-A/I-E, PE/Cyanine5 anti-mouse I-A/I-E, PE/Fire™ 744 anti-mouse I-A/I-E, Alexa Fluor® 488 anti-mouse I-A/I-E, Alexa Fluor® 647 anti-mouse I-A/I-E, Pacific Blue™ anti-mouse I-A/I-E, Biotin anti-mouse I-A/I-E, PE anti-mouse I-A/I-E, Purified anti-mouse I-A/I-E, PerCP/Cyanine5.5 anti-mouse I-A/I-E, PerCP anti-mouse I-A/I-E, Ultra-LEAF™ Purified anti-mouse I-A/I-E, TotalSeq™-B0117 anti-mouse I-A/I-E, Spark YG™ 593 anti-mouse I-A/I-E (Flexi-Fluor™) Antibody, FITC anti-mouse I-A/I-E, TotalSeq™-A0117 anti-mouse I-A/I-E, PE/Cyanine7 anti-mouse I-A/I-E, APC anti-mouse I-A/I-E, Spark PLUS B574™ anti-mouse I-A/I-E Antibody, Alexa Fluor® 700 anti-mouse I-A/I-E, TotalSeq™-C0117 anti-mouse I-A/I-E, Spark Blue™ 550 anti-mouse I-A/I-E, PE/Fire™ 640 anti-mouse I-A/I-E

Product Data



C57BL/6 mouse splenocytes were stained with anti-mouse/human CD45R/B220 (clone RA3-6B2) Spark PLUS UV395™ and anti-mouse I-A/I-E (clone M5/114.15.2) PE/Fire™ 744 (left) or rat IgG2b, κ PE/Fire™ 744 isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587