

PerCP/Fire™ 780 anti-human CD123 Antibody

Catalog# / Size	396731 / 25 tests 396732 / 100 tests
Clone	S18016F
Regulatory Status	RUO
Other Names	IL-3R α , IL-3 Receptor alpha
Isotype	Mouse IgG1, κ
Description	CD123 is the 70 kD transmembrane α chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common β chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the β chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Hu CD123 transfectants
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PerCP/Fire™ 780 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP/Fire™ 780 has a maximum excitation of 478 nm and a maximum emission of 780 nm.
Excitation Laser	Blue Laser (488 nm)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, associates with CDw131, 70 kD.
Distribution	Myeloid precursors, basophils, mast cells, macrophages, dendritic cells, megakaryocytes, subset of lymphocytes.
Function	Hematopoietic cell proliferation, differentiation
Ligand/Receptor	IL-3
Cell Type	Basophils, Dendritic cells, Hematopoietic stem and progenitors, Lymphocytes, Macrophages, Mast

cells, Megakaryocytes

Biology Area	Immunology
Molecular Family	CD Molecules, Cytokine/Chemokine Receptors
Antigen References	1. Miyajima A, <i>et al.</i> 1993. <i>Blood</i> . 82:1960.
Gene ID	3563


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-human CD123, Purified anti-human CD123, KIRAVIA Blue 520™ anti-human CD123, PE/Dazzle™ 594 anti-human CD123, APC anti-human CD123, APC/Fire™ 750 anti-human CD123 Antibody, PE/Cyanine7 anti-human CD123 Antibody, PerCP/Cyanine5.5 anti-human CD123 Antibody, Spark NIR™ 685 anti-human CD123, Spark Violet™ 423 anti-human CD123 Antibody, Biotin anti-human CD123, PE/Fire™ 810 anti-human CD123, Spark UV™ 387 anti-human CD123, Spark PLUS UV395™ anti-human CD123, PerCP/Fire™ 780 anti-human CD123 Antibody

Product Data



Human peripheral blood lymphocytes were stained with anti-human CD45RO (clone UCHL1) Brilliant Violet 421™ and anti-human CD123 (clone S18016F) PerCP/Fire™ 780 (left) or mouse IgG1, κ PerCP/Fire™ 780 isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587