

PerCP/Fire™ 780 anti-human CD28 Antibody

Catalog# / Size	377817 / 25 tests 377818 / 100 tests
Clone	S20013B
Regulatory Status	RUO
Other Names	T44, Tp44
Isotype	Mouse IgG1, κ
Description	CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. <i>In vitro</i> studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell activation and proliferation.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human CD28 transfected cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PerCP/Fire™ 780 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP/Fire™ 780 has a maximum excitation of 478 nm and a maximum emission of 780 nm.
Excitation Laser	Blue Laser (488 nm)
Application Notes	Clone S20013B antibody can completely block the binding of clone CD28.2 on target. Clone S20013B can partially block Clone S20013F on target. Clone S20013B can increase anti-CD3 induced human peripheral T-cell proliferation. Clone S20013B is a superagonistic antibody raised towards human CD28, and can induce activation, proliferation of human peripheral T cells without anti-CD3 stimulation.
RRID	AB_3683416 (BioLegend Cat. No. 377817) AB_3683416 (BioLegend Cat. No. 377818)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, homodimer, 44 kD
Distribution	Mature T cells, thymocytes, NK cell subsets, plasma cells, EBV-positive B cells
Function	T cell costimulation
Ligand/Receptor	CD80, CD86
Cell Type	Lymphocytes, T cells, Tfh, Th1, Th17, Th2, Thymocytes, Tregs
Biology Area	Adaptive Immunity, Cell Proliferation and Viability, Costimulatory Molecules, Immuno-Oncology, Immunology
Molecular Family	CD Molecules

Antigen References

- Schlossman S, *et al.* Eds. 1995. Leukocyte Typing V. Oxford University Press. New York.
- June CH, *et al.* 1994. *Immunol Today*. 15:321-31.
- Linsley PS and Ledbetter JA. 1993. *Annu Rev Immunol*. 11:191-212.

Gene ID [940](#)


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Ultra-LEAF™ Purified anti-human CD28 (Superagonistic), APC anti-human CD28, FITC anti-human CD28, PE anti-human CD28, PE/Cyanine7 anti-human CD28, PerCP/Cyanine5.5 anti-human CD28, PerCP/Fire™ 806 anti-human CD28, PerCP/Fire™ 780 anti-human CD28 Antibody, Spark Blue™ 550 anti-human CD28 (Flexi-Fluor™), Spark Blue™ 574 anti-human CD28 (Flexi-Fluor™) Antibody

Product Data



Human peripheral blood lymphocytes were stained with anti-human CD3 (clone UCHT1) FITC and anti-human CD28 (clone S20013B) PerCP/Fire™ 780 (left) or mouse IgG1, κ PerCP/Fire™ 780 isotype control (right)

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587