

## PerCP/Fire™ 806 anti-human CD183 (CXCR3) Antibody

<b>Catalog# / Size</b>	353771 / 25 tests 353772 / 100 tests
<b>Clone</b>	G025H7
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	CXCR3, G protein-coupled receptor 9 (GPR9), CKR-L2
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	Human CXCR3, also known as GPR9, is a chemokine receptor that binds CXCL9, CXCL10, and CXCL11. It is a 38 kD seven-pass transmembrane receptor coupled to G-protein. CXCR3 is highly expressed by T cells (Th1), natural killer cells (NK cells), dendritic cells, mast cells, alveolar macrophages, eosinophils, and human airway epithelial cells. CXCR3 is important for effector lymphocyte recruitment into inflamed tissue in various inflammatory and autoimmune diseases, such as chronically inflamed liver, Crohn's disease, rheumatoid arthritis, multiple sclerosis, and inflammatory skin diseases.

### Product Details

---

<b>Verified Reactivity</b>	Human, Cynomolgus, Rhesus
<b>Reported Reactivity</b>	African Green, Baboon
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Human CXCR3 transfectants
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with PerCP/Fire™ 806 under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our <a href="#">Certificate of Analysis</a> online tool.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.  * PerCP/Fire™ 806 has a maximum excitation of 478 nm and a maximum emission of 806 nm.
<b>Excitation Laser</b>	Blue Laser (488 nm)
<b>RRID</b>	AB_3675076 (BioLegend Cat. No. 353771) AB_3675076 (BioLegend Cat. No. 353772)

### Antigen Details

---

<b>Structure</b>	CXC-chemokine receptor, G protein-coupled receptor, seven-pass transmembrane receptor
<b>Distribution</b>	T cell subset, NK cells, plasmacytoid dendritic cells, GM-CSF activated CD34 <sup>+</sup> hematopoietic

progenitors, mast cells, alveolar macrophages, eosinophils, and airway epithelial cells

<b>Function</b>	Essential in T cell recruitment to sites of inflammation
<b>Ligand/Receptor</b>	CXCL9, CXCL10, and CXCL11
<b>Cell Type</b>	Dendritic cells, Eosinophils, Epithelial cells, Hematopoietic stem and progenitors, Macrophages, Mast cells, NK cells, T cells, Tregs
<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	CD Molecules, Cytokine/Chemokine Receptors, GPCR
<b>Antigen References</b>	1. Loetscher M, <i>et al.</i> 1996. <i>J. Exp. Med.</i> 184:963. 2. Cole KE, <i>et al.</i> 1998. <i>J. Exp. Med.</i> 187:2009. 3. Aksoy MO, <i>et al.</i> 2006. <i>Am. J. Physiol. Lung Cell Mol. Physiol.</i> 290:L909. 4. Curbishley SM, <i>et al.</i> 2005. <i>Am. J. Pathol.</i> 167:887. 5. Turner JE, <i>et al.</i> 2007. <i>Mini. Rev. Med. Chem.</i> 7:1089. 6. Wenzel J, <i>et al.</i> 2008. <i>J. Invest. Dermatol.</i> 128:67.
<b>Gene ID</b>	<a href="#">2833</a>


## Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-human CD183 (CXCR3), APC/Cyanine7 anti-human CD183 (CXCR3), FITC anti-human CD183 (CXCR3), PE anti-human CD183 (CXCR3), APC anti-human CD183 (CXCR3), Alexa Fluor® 488 anti-human CD183 (CXCR3), Alexa Fluor® 647 anti-human CD183 (CXCR3), PerCP/Cyanine5.5 anti-human CD183 (CXCR3), Brilliant Violet 421™ anti-human CD183 (CXCR3), PE/Cyanine7 anti-human CD183 (CXCR3), Pacific Blue™ anti-human CD183 (CXCR3), Brilliant Violet 510™ anti-human CD183 (CXCR3), Brilliant Violet 605™ anti-human CD183 (CXCR3), Brilliant Violet 650™ anti-human CD183 (CXCR3), Brilliant Violet 711™ anti-human CD183 (CXCR3), Purified anti-human CD183 (CXCR3) (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD183 (CXCR3), PerCP anti-human CD183 (CXCR3), Brilliant Violet 785™ anti-human CD183 (CXCR3), Alexa Fluor® 700 anti-human CD183 (CXCR3), Biotin anti-human CD183 (CXCR3), TotalSeq™-A0140 anti-human CD183 (CXCR3), TotalSeq™-C0140 anti-human CD183 (CXCR3), Ultra-LEAF™ Purified anti-human CD183 (CXCR3), TotalSeq™-B0140 anti-human CD183 (CXCR3), APC/Fire™ 750 anti-human CD183 (CXCR3), TotalSeq™-D0140 anti-human CD183 (CXCR3), PE/Fire™ 810 anti-human CD183 (CXCR3) Antibody, APC/Fire™ 810 anti-human CD183 (CXCR3) Antibody, PE/Cyanine5 anti-human CD183 (CXCR3) Antibody, PE/Fire™ 640 anti-human CD183 (CXCR3), PerCP/Fire™ 780 anti-human CD183 (CXCR3), Brilliant Violet 570™ anti-human CD183 (CXCR3), Spark Red™ 718 anti-human CD183 (CXCR3) (Flexi-Fluor™), PE/Fire™ 700 anti-human CD183 (CXCR3), PerCP/Fire™ 806 anti-human CD183 (CXCR3), PE anti-human CD183

## Product Data



Human peripheral blood lymphocytes were stained with anti-human CD3 (clone UCHT1) Brilliant Violet 421™ and anti-human CD183 (CXCR3) (clone G025H7) PerCP/Fire™ 806 (left) or mouse IgG1, κ PerCP/Fire™ 806 isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at [www.biolegend.com/terms](http://www.biolegend.com/terms) ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of

BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587