

Spark PLUS B488™ anti-human CD274 (B7-H1, PD-L1) Antibody

Catalog# / Size	329763 / 25 tests 329764 / 100 tests
Clone	29E.2A3
Regulatory Status	RUO
Other Names	Programmed cell death ligand 1 (PD-L1), B7 homolog 1 (B7-H1)
Isotype	Mouse IgG2b, κ
Description	CD274, also known as PD-L1 and B7-H1, is type I transmembrane glycoprotein that serves as a ligand for CD279 (PD-1). This interaction is believed to regulate the balance between the stimulatory and inhibitory signals needed for responses to microbes and maintenance of self-tolerance. CD274 is involved in the costimulation of T cell proliferation and IL-10 and IFN- γ production in an IL-2-dependent and CD279-independent manner. Conflicting data has shown that CD274 can inhibit T cell proliferation and cytokine production, and alternatively, enhance T cell activation. Other studies suggest that CD274 may signal bidirectionally, raising interesting implications for its expression in a wide variety of cell types, including T and B cells, antigen-presenting cells, and nonhematopoietic cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Full length human PD-L1
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS B488™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS B488™ has a maximum excitation of 499 nm and a maximum emission of 520 nm.
Excitation Laser	Blue Laser (488 nm)
Application Notes	Clone 29E.2A3 is reported to recognize an epitope on PD-L1 within the PD-L1-CD80 binding region ⁵ . Additional reported applications (for the relevant formats) include: blocking ¹⁻³ and immunohistochemical staining of acetone-fixed frozen sections ¹ . The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 329715, 329716, 329745 - 329748).

It has been observed that clone 29E.2A3 is able to bind to Alexa Fluor® 700 antibody conjugates during multi-color immunofluorescent staining. This interaction can be resolved by sequentially

staining with the 29E.2A3 antibody first and then followed by the Alexa Fluor® 700 conjugate of interest.

Clone 29E.2A3 does not work in Western blot applications⁷.

Application References

(PubMed link indicates
BioLegend citation)

1. Brown J, *et al.* 2003. *J. Immunol.* 170:1257. (FC, IHC, Block)
2. Radziejewicz H, *et al.* 2007. *J. Virol.* 81:2545. (Block)
3. Nakamoto N, *et al.* 2009. *PLoS Pathog.* 5:e1000313. (Block)
4. Barsoum IB, *et al.* 2014. *Cancer Res.* 74:665. [PubMed](#)
5. Haile, S *et al.* 2013. *J. Immunol.* 191:2829.
6. RL M, *et al.* 2015. *PNAS.* 112:6506-6514. [PubMed](#)
7. Mahoney KM, *et al.* 2015. *Cancer Immunol. Res.* 3:1308.

RRID

AB_3720643 (BioLegend Cat. No. 329763)

AB_3720643 (BioLegend Cat. No. 329764)

Antigen Details

Distribution	T cells, B cells, NK cells, monocytes/macrophages, granulocytes and dendritic cells
Function	CD274 is involved in the costimulatory signal, essential for T lymphocyte proliferation and production of IL-10 and IFN- γ , in an IL-2-dependent and a PD-1-CD1-independent manner. Its interaction with PD-1-CD1 inhibits T-cell proliferation and cytokine production.
Ligand/Receptor	PD-1 (PDCD1)
Cell Type	B cells, Dendritic cells, Fibroblasts, Granulocytes, Macrophages, Monocytes, NK cells, T cells
Biology Area	Cancer Biomarkers, Costimulatory Molecules, Immunology
Molecular Family	Adhesion Molecules, CD Molecules, Immune Checkpoint Receptors
Antigen References	1. Sharpe A, <i>et al.</i> 2007. <i>Nat. Immunol.</i> 8:239.
Gene ID	29126


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

TotalSeq™-D0007 anti-human CD274 (B7-H1, PD-L1), Spark YG™ 570 anti-human CD274 (B7-H1, PD-L1), PE/Cyanine5 anti-human CD274 (B7-H1, PD-L1), PE/Fire™ 810 anti-human CD274 (B7-H1, PD-L1) Antibody, Purified anti-human CD274 (B7-H1, PD-L1) (Maxpar® Ready), GolnVivo™ Purified anti-human CD274 (B7-H1, PD-L1), PE/Dazzle™ 594 anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 605™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 650™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 510™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 785™ anti-human CD274 (B7-H1, PD-L1), PerCP/Cyanine5.5 anti-human CD274 (B7-H1, PD-L1), Spark PLUS UV395™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 711™ anti-human CD274 (B7-H1, PD-L1), Purified anti-human CD274 (B7-H1, PD-L1), TotalSeq™-A0007 anti-human CD274 (B7-H1, PD-L1), TotalSeq™-C0007 anti-human CD274 (B7-H1, PD-L1), Alexa Fluor® 594 anti-human CD274 (B7-H1, PD-L1), Spark PLUS B488™ anti-human CD274 (B7-H1, PD-L1), Biotin anti-human CD274 (B7-H1, PD-L1), PE anti-human CD274 (B7-H1, PD-L1), TotalSeq™-B0007 anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 421™ anti-human CD274 (B7-H1, PD-L1), Ultra-LEAF™ Purified anti-human CD274 (B7-H1, PD-L1), PE/Cyanine7 anti-human CD274 (B7-H1, PD-L1), APC anti-human CD274 (B7-H1, PD-L1)

Product Data



PHA-stimulated (3 days) human peripheral blood mononuclear cells were stained with anti-human CD3 (clone UCHT1) APC and with anti-human CD274 (B7-H1, PD-L1) (clone 29E.2A3) Spark PLUS B488™ (left) or stained with anti-human CD3 (clone UCHT1) APC only (right). Data shown is gated on the lymphocyte population.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587