

Spark PLUS UV395™ anti-human CD117 (c-kit) Antibody

Catalog# / Size	375217 / 25 tests 375218 / 100 tests
Clone	S18022G
Regulatory Status	RUO
Other Names	Stem cell factor receptor, c-kit, mast cell growth factor receptor, steel factor receptor
Isotype	Mouse IgG2a, κ
Description	CD117 is a 145 kD protein tyrosine kinase also known as c-kit. It is a receptor for stem cell factor or c-kit ligand. CD117 is expressed on pluripotent hematopoietic progenitor cells (approximately 1-4% bone marrow cells), mast cells, and acute myeloid leukemia cells (AML). CD117 binding of c-kit ligand induces phosphorylation of CD117 and stimulates proliferation and survival of primitive hematopoietic stem cells as well as erythroid-committed and granulomonocytic committed cells.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human CD117 transfectant
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS UV395™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS UV395™ has a maximum excitation of 355 nm and a maximum emission of 385 nm.
Excitation Laser	Ultraviolet Laser (355 nm)
Application Notes	Clone S18022G antibody can completely block the binding of clone 104D2 to the target, but it does not block the binding of clone w18195C to the target. Clone S18022G antibody binding to the target is not affected after cells are fixed by 4% PFA.
RRID	AB_3699015 (BioLegend Cat. No. 375217) AB_3699015 (BioLegend Cat. No. 375218)

Antigen Details

Structure	Growth factor receptor with tyrosine kinase activity, subclass III, approximately 145 kD
------------------	--

Distribution	Pluripotent hematopoietic progenitor cells (approximately 1-4% bone marrow cells), mast cells, acute myeloid leukemic cells (AML)
Function	Receptor for stem cell factor. Induces proliferation and survival of primitive hematopoietic progenitors. Potent inducer of proliferation in erythroid-committed progenitor cells. Defects in CD117 have been linked to severe anemia and a decreased number of hematopoietic progenitor cells.
Ligand/Receptor	Stem cell factor (c-kit ligand)
Antigen References	1. Giebel LB, <i>et al.</i> 1992. <i>Oncogene</i> 7:2207. 2. Furitsu T, <i>et al.</i> 1993. <i>J. Clin. Invest.</i> 92:1736.
Gene ID	3815


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Pacific Blue™ anti-human CD117 (c-kit) Antibody, PE/Cyanine7 anti-human CD117 (c-kit) Antibody, APC anti-human CD117 (c-kit), Purified anti-human CD117 (c-kit) Antibody, PE anti-human CD117 (c-kit) Antibody, Brilliant Violet 421™ anti-human CD117 (c-kit) Antibody, Brilliant Violet 570™ anti-human CD117 (c-kit) Antibody, Brilliant Violet 785™ anti-human CD117 (c-kit) Antibody, Spark PLUS UV395™ anti-human CD117 (c-kit)

Product Data



Kasumi-3 cells (human acute myeloblastic leukemia cell line) were stained with anti-human CD117 (c-kit) (clone S18022G) Spark PLUS UV395™ (filled histogram). Open histogram represents unstained cells.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587