

Spark PLUS UV395™ anti-human CD80 Antibody

Catalog# / Size	375415 / 25 tests 375416 / 100 tests
Clone	W17149D
Regulatory Status	RUO
Other Names	B7-1, B7, BB1
Isotype	Rat IgG2a, κ
Description	CD80, also known as B7-1, B7, and BB1, is a 60 kD single chain type I glycoprotein belonging to the immunoglobulin superfamily. CD80 is expressed on activated B and T cells, macrophages, and dendritic cells. CD80 binds to CD28 and CD152 (CTLA-4). Along with CD86, CD80 plays a critical role in regulation of T cell activation. The interaction of CD80 with CD28 provides a potent costimulatory signal for T cell activation through the CD3 complex, while its interaction with CTLA-4 provides an inhibitory signal for T cell activation.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	RBL1 cells stably transfected with human CD80 (full length)
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS UV395™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS UV395™ has a maximum excitation of 355 nm and a maximum emission of 385 nm.
Excitation Laser	Ultraviolet Laser (355 nm)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, 60 kD
Distribution	Activated B cells and T cells, macrophages, and dendritic cells
Function	T cell costimulation
Ligand/Receptor	CD28, CD152 (CTLA-4)
Cell Type	B cells, Dendritic cells, Macrophages, T cells, Tregs

Biology Area	Cell Biology, Costimulatory Molecules, Immunology, Neuroscience, Neuroscience Cell Markers
Molecular Family	CD Molecules, Immune Checkpoint Receptors
Antigen References	<ol style="list-style-type: none"> 1. Lim TS, <i>et al.</i> 2012. PLoS One. 7(9): e45185. 2. Kamphorst, Alice O, <i>et al.</i> 2017. <i>Science</i>. 31;355(6332):1423-1427. 3. Zheng Y, <i>et al.</i> 2004. <i>J. Immunol.</i> 1;172(5):2778-84.
Gene ID	941


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD80 Antibody, APC anti-human CD80 Antibody, FITC anti-human CD80 Antibody, PE/Cyanine7 anti-human CD80, PE anti-human CD80, PerCP/Cyanine5.5 anti-human CD80, PE/Fire™ 744 anti-human CD80, Spark Blue™ 574 anti-human CD80 (Flexi-Fluor™), Spark Blue™ 550 anti-human CD80 (Flexi-Fluor™), Spark PLUS UV395™ anti-human CD80 Antibody

Product Data



Raji cells (human B-cell Burkitts lymphoma cell line) were stained with anti-human CD80 (clone W17149D) Spark PLUS UV395™ (filled histogram) or rat IgG2a, κ Spark PLUS UV395™ isotype control (open histogram).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587