

Spark PLUS UV395™ anti-human HLA-DR, DP, DQ Antibody

Catalog# / Size	361727 / 25 tests 361728 / 100 tests
Clone	Tü39
Regulatory Status	RUO
Other Names	MHC class II, Major Histocompatibility complex II, human leukocyte antigen, HLA
Isotype	Mouse IgG2a, κ
Description	HLA-DR, HLA-DP, and HLA-DQ are heterodimeric cell surface glycoproteins comprised of an α (heavy) chain and a β (light) chain. They are expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. Variations in the HLA gene expression are crucial to graft survival.

Product Details

Verified Reactivity	Human
Reported Reactivity	Baboon, Cynomolgus, Cat, Dog, Cow
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human PBL
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS UV395™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS UV395™ has a maximum excitation of 355 nm and a maximum emission of 385 nm.
Excitation Laser	Ultraviolet Laser (355 nm)
Application Notes	Tü39 has been reported to react with a shared epitope of HLA-DR, HLA-DP, and HLA-DQ. Additional reported applications (of relevant formats) include immunoprecipitation ⁶ , <i>in vitro</i> blocking of MLR ⁵ , and suppressor cell generation ⁴ . The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (contact our custom solutions team).

Application References

(PubMed link indicates
BioLegend citation)

1. Pawelec G, *et al.* 1985. *Hum. Immunol.* 12:165.
2. Shaw S, *et al.* 1985. *Hum. Immunol.* 12:191.
3. Ziegler A, *et al.* 1986. *Immunobiology* 171:77.
4. Pawelec G, *et al.* 1986. *Hum. Immunol.* 17:343. (Block)
5. Dai Z, *et al.* 2009. *J. Exp. Med.* 206:793. (Block)
6. Pawelec G, *et al.* 1988. *J. Exp. Med.* 167:243. (IP)

RRID

AB_3699004 (BioLegend Cat. No. 361727)
AB_3699004 (BioLegend Cat. No. 361728)

Antigen Details

Structure	Ig superfamily, MHC class II, heterodimeric transmembrane protein
Distribution	B cells, activated T cells, monocytes/macrophages, dendritic cells, other APCs
Function	Antigen presentation
Ligand/Receptor	CD3/TCR, CD4
Cell Type	Antigen-presenting cells, B cells, Dendritic cells, Macrophages, Monocytes, T cells
Biology Area	Immunology, Innate Immunity
Molecular Family	MHC Antigens
Antigen References	

1. Thorsby E. 1974. *Transplant. Rev.* 18:51.
2. Qvigstad E, *et al.* 1984. *Hum. Immunol.* 11:207.
3. Serenius B, *et al.* 1984. *EMBO J.* 3:3209.
4. Ottenhoff TH, *et al.* 1985. *Hum. Immunol.* 13:105.
5. Strassmann G, *et al.* 1985. *Hum. Immunol.* 13:125.
6. Trowsdale J, *et al.* 1985. *Immunol Rev.* 85:5.

Gene ID

[3122](#)
[3123](#)
[3113](#)
[3115](#)
[3117](#)
[3119](#)


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human HLA-DR, DP, DQ, FITC anti-human HLA-DR, DP, DQ, Alexa Fluor® 647 anti-human HLA-DR, DP, DQ, PE/Cyanine7 anti-human HLA-DR, DP, DQ, APC anti-human HLA-DR, DP, DQ, PerCP/Cyanine5.5 anti-human HLA-DR, DP, DQ, PE anti-human HLA-DR, DP, DQ, APC/Fire™ 750 anti-human HLA-DR, DP, DQ, TotalSeq™-A1018 anti-human HLA-DR, DP, DQ, TotalSeq™-D1018 anti-human HLA-DR, DP, DQ, TotalSeq™-B1018 anti-human HLA-DR, DP, DQ, Spark Blue™ 574 anti-human HLA-DR, DP, DQ, TotalSeq™-C1018 anti-human HLA-DR, DP, DQ, Spark Red™ 718 anti-human HLA-DR/HLA-DP/HLA-DQ (Flexi-Fluor™), Spark Blue™ 550 anti-human HLA-DR/HLA-DP/HLA-DQ (Flexi-Fluor™), Spark PLUS UV395™ anti-human HLA-DR, DP, DQ, PE/Fire™ 700 anti-human HLA-DR, DP, DQ Antibody

Product Data



Human peripheral blood lymphocytes were stained with anti-human CD19 (clone HIB19) APC and anti-human HLA-DR, DP, DQ (clone Tü39) Spark PLUS UV395™ (left) or mouse IgG2a, κ Spark PLUS UV395™ isotype control (right).

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587