

Spark PLUS YG581™ anti-human CD20 Antibody

Catalog# / Size	302385 / 25 tests 302386 / 100 tests
Clone	2H7
Regulatory Status	RUO
Workshop	IV B201
Other Names	B1, Bp35
Isotype	Mouse IgG2b, κ
Description	CD20 is a 33-37 kD, four transmembrane spanning protein, also known as B1 and Bp35. CD20 is expressed on pre-B-cells, resting and activated B cells (not plasma cells), some follicular dendritic cells, and at low levels on a T cell subset. CD20 is heavily phosphorylated on activated B cells and malignant B cells. Homo-oligomeric complexes of CD20 are thought to form Ca^{2+} conductive ion channels in the plasma membrane of B cells. The CD20 molecule is involved in B-cell activation and is associated with various Src family kinases (Lyn, Lck, Fyn). It exists in a complex with MHC class I and II, CD53, CD81, and CD82.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Reported Reactivity	Baboon, Capuchin Monkey, Chimpanzee, Pigtailed Macaque, Squirrel Monkey
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human tonsillar B cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS YG581™ under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration and expiration, please enter the lot number in our Certificate of Analysis online tool.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS YG581™ has a maximum excitation of 562 nm and a maximum emission of 581 nm.
Excitation Laser	Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	The epitope recognized by clone 2H7 has been mapped to the sequence YNCEPANPSEKNSPST which lies in the large extracellular loop of human CD20. Additional reported applications (for the relevant formats) include: immunoprecipitation ⁴ and immunohistochemical staining of acetone-fixed frozen sections ⁵ .

Application References

(PubMed link indicates
BioLegend citation)

1. Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
2. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
3. McMichael A, *et al.* Eds. 1987. Leucocyte Typing III Oxford University Press. New York.
4. Polyak MJ, *et al.* 2002. *Blood* 99:3256. (IP)
5. Mack CL, *et al.* 2004. *Pediatr. Res.* 56:79. (IHC)

Antigen Details

Structure	Four transmembrane protein (TM4SF), heavily phosphorylated after activation, 33-37 kD
Distribution	B cell, T cell subsets
Function	B cell activation
Ligand/Receptor	Src family tyrosine kinases, MHC class I, II, CD53, CD81, CD82
Cell Type	B cells, T cells
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	1. Hultin L, <i>et al.</i> 1993. <i>Cytometry</i> 14:196. 2. Tedder T, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:450.
Gene ID	931


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

TotalSeq™-D0100 anti-human CD20, GMP FITC anti-human CD20, GMP APC anti-human CD20, Spark Violet™ 500 anti-human CD20, GMP PE/Cyanine7 anti-human CD20, Spark Violet™ 538 anti-human CD20, Spark Blue™ 515 anti-human CD20, GMP PerCP/Cyanine5.5 anti-human CD20, PE anti-human CD20, Spark YG™ 593 anti-human CD20, GMP APC/Fire™ 750 anti-human CD20, GMP PE anti-human CD20, PE/Dazzle™ 594 anti-human CD20, FITC anti-human CD20, Alexa Fluor® 594 anti-human CD20, APC/Fire™ 750 anti-human CD20, Pacific Blue™ anti-human CD20, APC anti-human CD20, Purified anti-human CD20, PE/Cyanine5 anti-human CD20, Brilliant Violet 650™ anti-human CD20, Brilliant Violet 785™ anti-human CD20, Brilliant Violet 510™ anti-human CD20, Brilliant Violet 421™ anti-human CD20, Brilliant Violet 711™ anti-human CD20, Purified anti-human CD20 (Maxpar® Ready), GMP Pacific Blue™ anti-human CD20, Brilliant Violet 750™ anti-human CD20, Spark YG™ 581 anti-human CD20 (Flexi-Fluor™), Spark Blue™ 550 anti-human CD20 (Flexi-Fluor™), Spark Red™ 718 anti-human CD20 (Flexi-Fluor™), Biotin anti-human CD20, StarBright UltraViolet 665 anti-human CD20, Alexa Fluor® 488 anti-human CD20, TotalSeq™-A0100 anti-human CD20, TotalSeq™-B0100 anti-human CD20, TotalSeq™-C0100 anti-human CD20, APC/Fire™ 810 anti-human CD20, Spark Blue™ 574 anti-human CD20, Spark PLUS YG581™ anti-human CD20, Alexa Fluor® 647 anti-human CD20, Alexa Fluor® 700 anti-human CD20, PE/Cyanine7 anti-human CD20, PerCP/Cyanine5.5 anti-human CD20, APC/Cyanine7 anti-human CD20, Spark NIR™ 685 anti-human CD20, Brilliant Violet 570™ anti-human CD20, Brilliant Violet 605™ anti-human CD20, Alexa Fluor® 660 anti-human CD20, PerCP anti-human CD20

Product Data



Human peripheral blood cells were stained with anti-human CD19 (clone HIB19) PE/Fire™ 640 and anti-human CD20 (clone 2H7) Spark PLUS YG581™ (left) or stained with anti-human CD19 (clone HIB19) PE/Fire™ 640 only (right). Data shown were gated on the lymphocyte population.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587