

Spark PLUS YG581™ anti-mouse Ly-6G Antibody

Catalog# / Size	127687 / 25 µg
Clone	1A8
Regulatory Status	RUO
Other Names	Lymphocyte antigen 6 complex, locus G
Isotype	Rat IgG2a, κ
Description	Lymphocyte antigen 6 complex, locus G (Ly-6G), a 21-25 kD GPI-anchored protein, is expressed on the majority of myeloid cells in bone marrow and peripheral granulocytes.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Ly-6G transfected EL-4J cell line.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Preparation	The antibody was purified by affinity chromatography and conjugated with Spark PLUS YG581™ under optimal conditions.
Concentration	0.2 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application. * Spark PLUS YG581™ has a maximum excitation of 562 nm and a maximum emission of 581 nm.
Excitation Laser	Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	While 1A8 recognizes only Ly-6G, clone RB6-8C5 recognizes both Ly-6G and Ly-6C. Clone RB6-8C5 binds with high affinity to mouse Ly-6G molecules and to a lower extent to Ly-6C¹⁵. Clone RB6-8C5 impairs the binding of anti-mouse Ly-6G clone 1A8¹⁵. However, clone RB6-8C5 is able to stain in the presence of anti-mouse Ly-6C clone HK1.4¹⁶. Additional reported applications (for the relevant formats) include: immunohistochemistry⁹ of frozen sections¹⁰ and paraffin-embedded sections¹¹, depletion^{4, 12-14}, and spatial biology (IBEX)^{20,21}. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for <i>in vivo</i> studies or highly sensitive assays (Cat. No. 127632, 127649, 127650, 127661 and 127662).

Application References

(PubMed link indicates
BioLegend citation)

1. Fleming TJ, *et al.* 1993. *J. Immunol.* 151:2399. (FC)
2. Daley JM, *et al.* 2008. *J. Leukocyte Biol.* 83:1. (FC)
3. Dietlin TA, *et al.* 2007. *J. Leukocyte Biol.* 81:1205. (FC)
4. Daley J, *et al.* 2007. *J. Leukocyte Biol.* doi:10.1189. (Deplete) [PubMed](#)
5. Tadagavadi RK, *et al.* 2010. *J. Immunol.* 185:4904. [PubMed](#)
6. Sumagin R, *et al.* 2010. *J. Immunol.* 185:7057. [PubMed](#)
7. Guiducci C, *et al.* 2010. *J. Exp Med.* 207:2931. [PubMed](#)
8. Fujita M, *et al.* 2011. *Cancer Res.* 71:2664. [PubMed](#)
9. Van Leeuwen, *et al.* 2008. *Arterioscler. Thromb. Vasc. Biol.* 28:84. (IHC)
10. Kowanzet M, *et al.* 2010. *P. Natl. Acad. Sci. USA* 107:21248. [supplementary data] (IHC)
11. Esbona K, *et al.* 2016. *Breast Cancer Res.* 18:35. (IHC)
12. Wojtasiak M, *et al.* 2010. *J. Gen. Virol.* 91:2158. (FC, Deplete)

[See More](#)

Antigen Details

Structure	A 21-35 kD GPI-anchored membrane protein
Distribution	Expressed on the majority of myeloid cells in bone marrow and peripheral granulocytes. The monoclonal antibody RB6-8C5 recognizes both Ly-6G and Ly-6C.
Cell Type	Granulocytes, Macrophages, Monocytes
Biology Area	Immunology, Innate Immunity
Antigen References	Fleming TJ, <i>et al.</i> 1993. <i>J. Immunol.</i> 151:2399.
Gene ID	546644


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC/Fire™ 810 anti-mouse Ly-6G Antibody, PE/Cyanine5 anti-mouse Ly-6G, Spark YG™ 593 anti-mouse Ly-6G, PE/Fire™ 640 anti-mouse Ly-6G, Spark UV™ 387 anti-mouse Ly-6G, Spark YG™ 570 anti-mouse Ly-6G, Purified anti-mouse Ly-6G (Maxpar® Ready), PE/Fire™ 810 anti-mouse Ly-6G Antibody, Brilliant Violet 785™ anti-mouse Ly-6G, Brilliant Violet 711™ anti-mouse Ly-6G, PerCP anti-mouse Ly-6G, Brilliant Violet 605™ anti-mouse Ly-6G, PE/Dazzle™ 594 anti-mouse Ly-6G, APC/Fire™ 750 anti-mouse Ly-6G, Purified anti-mouse Ly-6G, FITC anti-mouse Ly-6G, PE anti-mouse Ly-6G, Alexa Fluor® 647 anti-mouse Ly-6G, Biotin anti-mouse Ly-6G, Brilliant Violet 570™ anti-mouse Ly-6G, Alexa Fluor® 488 anti-mouse Ly-6G, PE/Cyanine7 anti-mouse Ly-6G, Ultra-LEAF™ Purified anti-mouse Ly-6G, Brilliant Violet 421™ anti-mouse Ly-6G, Spark Red™ 718 anti-mouse Ly-6G (Flexi-Fluor™), Spark Blue™ 574 anti-mouse Ly-6G (Flexi-Fluor™), Spark PLUS YG581™ anti-mouse Ly-6G, Brilliant Violet 650™ anti-mouse Ly-6G, PerCP/Fire™ 806 anti-mouse Ly-6G, StarBright UltraViolet 575 anti-mouse Ly-6G, Spark NIR™ 685 anti-mouse Ly-6G, Pacific Blue™ anti-mouse Ly-6G, APC anti-mouse Ly-6G, PerCP/Cyanine5.5 anti-mouse Ly-6G, Alexa Fluor® 700 anti-mouse Ly-6G, APC/Cyanine7 anti-mouse Ly-6G, Brilliant Violet 510™ anti-mouse Ly-6G, Alexa Fluor® 594 anti-mouse Ly-6G, TotalSeq™-A0015 anti-mouse Ly-6G, TotalSeq™-C0015 anti-mouse Ly-6G, TotalSeq™-B0015 anti-mouse Ly-6G, Spark Blue™ 550 anti-mouse Ly-6G

Product Data



C57BL/6 mouse bone marrow cells were stained with anti-mouse/human CD11b (clone M1/70) FITC and anti-mouse Ly-6G (clone 1A8) Spark PLUS YG581™ (left) or stained with anti-mouse/human CD11b (clone M1/70) FITC only (right). Data shown are gated on the myeloid population.

For Research Use Only. Not for diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587